

(43) International Publication Date 30 November 2000 (30.11.2000)

PCT

(10) International Publication Number WO 00/71116 A1

(51) International Patent Classification⁷: C07D 207/34

A61K 31/40,

M., Dileep [IN/IN]; Vaishali Apartments, Pocket A-12, 3C, Kalkaji Extension, New Delhi 110 019, Maharashtra (IN).

(21) International Application Number: PCT/IB00/00014

(22) International Filing Date: 6 January 2000 (06.01,2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 775/DEL/99

25 May 1999 (25.05.1999) IN

(71) Applicant (for all designated States except US): RAN-BAXY LABORATORIES LIMITED [IN/IN]; 19 Nehru Place, New Delhi 110 019, Maharashtra (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KUMAR, Yatendra [IN/IN]; U-26/5, DLF Qutab Enclave, Phase - III, Gurgaon 122 001, Haryana (IN). THAPER, Rajesh, Kumar [IN/IN]; B-26, Sunshine Apartments, Block "C", Sushant Lok - I, Gurgaon 122 002, Haryana (IN). KUMAR, S.,

(81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,

LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

00, 03, 02, VN, 10, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TI, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

//71116 A1

(54) Title: PROCESS FOR THE PRODUCTION OF AMORPHOUS ATORVASTATIN CALCIUM

(57) Abstract: A process for the preparation of amorphous atorvastatin calcium and hydrates thereof which comprises: (a) dissolving crystalline atorvastatin calcium in a non-hydroxylic solvent; (b) adding a non-polar hydrocarbon anti-solvent or adding the dissolved atorvastatin to the non-polar anti-solvent to precipitate out atorvastatin calcium; and (c) removing the solvent by filtration to afford amorphous atorvastatin calcium.

PROCESS FOR THE PRODUCTION OF AMORPHOUS ATORVASTATIN CALCIUM

5

FIELD OF THE INVENTION

The present invention relates to a process for the production of amorphous atorvastatin calcium.

10

15

20

BACKGROUND OF THE INVENTION

Atorvastatin is chemically [R-(R*,R*)]-2-(4-fluoro-phenyl)-β dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino) carbonyl]-1H pyrrole-1-heptanoic acid. Atorvastatin calcium, a synthetic HMG-CoA reductase inhibitor, is used for the treatment of hyperlipidemia and hypercholesterolemia, both of which are risk factors for arteriosclerosis and coronary heart disease. Open dihydroxy carboxylic acid, lactone and various salt forms of atorvastatin have been synthesized.

United States Patent 5,273,995, describes that R-form of the ring opened acid form has surprising inhibition of the biosynthesis of cholesterol. Atorvastatin in its calcium salt form, i.e. $[R-(R^*,R^*)]-2-(4-fluoro-phenyl)-\beta$, δ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino) carbomyl]-1H-pyrrole-1-heptanoic acid calcium salt (2:1) having Formula 1:

5

10

15

20

is more suited to formulations and has been recommended as a drug.

United States patents 5,003,080; 5,097,045; 5,103,024; 5,124,482; 5,149,837; 5,248,793; 5,280,126; 5,342,952, which are herein incorporated by reference, describe various processes and key intermediates for preparing atorvastatin.

Atorvastatin calcium produced by the processes described in the above mentioned United States patents does not give amorphous atorvastatin consistently but gives a mixture of its crystalline and amorphous forms, which has unsuitable filtration and drying characteristics and are not suitable for large-scale production.

PCT application, WO 97/03959, discloses novel crystalline forms of atorvastatin calcium designated as Form I, Form II, and Form IV and method for their preparation which provide more favourable filtration and drying characteristics.

PCT application WO 97/03960 describes a procedure for converting the crystalline form of atorvastatin to the amorphous form. Process disclosed therein comprises dissolving crystalline form-I atorvastatin in a non-hydroxylic solvent like tetrahydrofuran or mixtures of tetrahydrofuran and toluene. The process involves complete removal of the solvent under high temperature (about 90°C) and high vacuum (about 5mm) using capital intensive equipment. Exposure of the material to high temperature for several days leads to degradation of the product. This makes the process very inconvenient to operate at a large scale. Slow removal of solvents at a manufacturing scale renders this process as inefficient cost-wise and less productive.

10

5

SUMMARY OF THE INVENTION

It is an objective of the present invention to provide an efficient process for the production of amorphous atorvastatin, which eliminates the problems of prior art and is convenient to operate on a commercial scale.

15

Accordingly, the present invention provides a process for the preparation of atorvastatin calcium in an amorphous form which comprises dissolving crystalline atorvastatin in a non-hydroxylic solvent, adding a suitable non-polar hydro-carbon solvent and recovering atorvastatin from a solution thereof, by solvent precipitation, isolating and drying the product.

20

Generally, the product can be isolated by any standard method known in the art such as by filtration, centrifugation or decantation. Typically, this product is isolated by filtration when any of the solvents within the scope of the process are used.

Major advantages of the present invention compared to the prior art processes are:

i. elimination of the need to remove solvent by drying techniques.

- ii. less time consuming with improved filtration.
- iii. easy to operate on large-scale.
- iv. reproducibly produces amorphous product having allowable levels of residual solvents.

The present invention thus provides a novel process for the preparation of amorphous atorvastatin calcium and hydrates thereof which comprises:

- (a) dissolving crystalline atorvastatin calcium in a non-hydroxylic solvent;
- (b) adding a non-polar hydrocarbon anti-solvent to precipitate out the material; and
- (c) removing the solvent by filtration to afford amorphous atorvastatin calcium

The non-hydroxylic solvent is selected from a group of solvents, which have the ability to dissolve crystalline atorvastatin and includes tetrahydrofuran. Suitable non-polar hydrocarbon solvents are selected from a group consisting of: n-hexane, n-heptane, cyclohexane, hexane fraction, heptane fraction or the like. In a preferred embodiment of this invention, the non-hydroxylic solvent is tetrahydrofuran and antisolvent is n-hexane, cyclohexane or n-heptane.

Generally, crystalline atorvastatin calcium is dissolved in a non-hydroxylic solvent, e.g. tetrahydrofuran, at a concentration of about 15% w/v to about 40% w/v, preferably at a concentration of about 25% w/v to about 15% w/v at ambient

5

15

20

temperature and a non-polar hydrocarbon, preferably n-hexane, cyclohexane or n-heptane, is added at 0°C to 50°C, preferably at 20°C to 25°C. The product is recovered by filtration at ambient temperature. Filtration, which is fast and smooth, is carried out using nutsche filtration or centrifuge filtration. Preferably, nutsche filtration is used on large scale preparation. Filtered material, a semi-dry powder, is further dried to remove surface solvents in a vacuum tray drier, tray drier, fluid bed drier or a rotary vacuum drier to afford amorphous material. Preferably, material is dried in a vacuum tray drier at about 20°C to about 80°C for 6 hours to 24 hours. Most preferably, drying is carried out at about 50°C to about 60°C for 12 hours.

10

5

Quantity of antisolvent varies from 5 times to 50 times the input of crystalline atorvastatin calcium depending upon its solution in non-hydroxylic solvent. Preferably, the quantity of antisolvent used is about 20 times to about 40 times the input of crystalline atorvastatin calcium to make overall concentration of about 5% w/v to about 2.5 w/v%.

15

20

Amorphous atorvastatin calcium prepared according to the process of the present invention may be characterized by its x-ray powder diffration pattern (Figures 2) as shown in the accompanied drawings. X-ray powder diffration patterns (Figures 2) show no peaks which are characteristic of a crystalline atorvastatin calcium (Figure 1 of the accompanied drawings) thus demonstrating the amorphous nature of the product.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is the diffractogram of crystalline atorvastatin calcium. The horizontal axis represents 2θ and the vertical axis corresponds to peak intensity.

Figure 2 is diffractogram of amorphous atorvastatin calcium. The horizontal axis represents 2θ and the vertical axis corresponds to peak intensity.

The present invention is illustrated by the following examples, which are not intended to limit the effective scope of the claims.

DETAILED DESCRIPTION OF THE INVENTION

Example 1

[R-(R*,R*)-2-(4-fluorophenyl)- β , δ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl)-1H-pyrrole-1-heptanoic acid hemicalcium salt (Amorphous Atorvasatin calcium).

Method A

5

10

15

20

Crystalline atorvastatin calcium (10 kg) was dissolved in tetrahydrofuran (30 lt) under stirring at ambient temperature. Clear solution so obtained was added slowly to cyclohexane (350 lt) under nitrogen atmosphere. It was vigorously stirred maintaining temperature at 20-25°C. The precipitated product was centrifuged and dried under vacuum at about 60°C for 12 hours. Atorvastatin (9.5 kg) in an

amorphous form was obtained having residual solvent levels of 0.01% w/w tetrahydrofuran and 0.6% w/w cyclohexane. X-ray powder diffraction pattern (Figure 2 as shown in the accompanied drawings) demonstrate the amorphous nature of the product.

5

10

Method B

Crystalline atorvastatin calcium (10 kg) was dissolved in tetrahydrofuran (30 lt) under stirring at ambient temperature. To a clear solution of atorvastatin, cyclohexane (350 lt) was added under vigorous stirring at 20 to 25°C. The precipitated mass was further stirred for 30 minutes and filtered in a centrifuge. The product was dried under vacuum at about 60°C for 12 hours. Atorvastatin (9.6 kg) in an amorphous form was obtained having residual solvent levels of 0.01% w/w for tetrahydrofuran and 0.7% w/w for cyclohexane. X-ray powder diffraction pattern demonstrates the amorphous nature of the product.

15

Example 2

 $[R-(R^*,R^*)-2-(4-fluorophenyl)\beta, \qquad \delta-dihydroxy-5-(1-methylethyl)-3-phenyl-4-\\ [(phenylamino)carbonyl)-1H-pyrrole-1-heptanoic acid hemicalcium salt (Amorphous Atorvasatin calcium)$

20

The process of Example 1 was repeated with crystalline atorvastatin calcium (10 kg) dissolved in tetrahydrofuran (30 lt) and using n-hexane instead of cyclohexane

to give amorphous atorvastatin (9.5 kg.). X-ray crystallography confirmed the amorphous nature of the product.

Example 3

5 [R-(R*,R*)]-2-4-fluorophenyl)-β, δ-dihydroxy-5-(1-methylethyl)-3-phenyl-4[(phenylamino)carbonyl)-1H-pyrrole-1-heptanoic acid hemicalcium salt
(Amorphous Atorvasatin calcium)

10

15

The process of Example 1 was repeated with crystalline atorvastatin calcium (10 kg) dissolved in tetrahydrofuran (30 lt) and using n-heptane instead of cylcohexane to give amorphous atorvastatin (9.6 kg). X-ray crystallography examination confirmed the amorphous nature of the product.


While the present invention has been described in terms of its specific embodiments, certain modifications and equivalents will be apparent to those skilled in the art and are intended to be included within the scope of the present invention.

WE CLAIM:

 A process for the preparation of amorphous atorvastatin calcium and hydrates thereof which comprises:


- (a) dissolving crystalline atorvastatin calcium in a non-hydroxylic solvent;
- (b) adding a non-polar hydrocarbon anti-solvent or adding the dissolved atorvastatin to the non-polar anti-solvent to precipitate out atorvastatin calcium; and
- (c) removing the solvent by filtration to afford amorphous atorvastatin calcium.
- 2. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is chosen from a group of non-polar hydrocarbon solvents comprising n-hexane, cyclohexane or n-heptane.
- 3. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is n-hexane.
- 4. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is cylcohexane.
- 5. The process of claim 1, wherein the non-hydroxylic solvent is tetrahydrofuran and anti-solvent is n-heptane.
- 6. The process of claim 1, wherein said amorphous atorvastatin calcium is isolated by filtration.





WO 00/71116





Interr nal Application No PCT/IB 00/00014

A CLASSIF IPC 7	FICATION OF SUBJECT MATTER A61K31/40 C07D2O7/34		
According to	o International Patent Classification (IPC) or to both national classif	lication and IPC	
B. FIELDS S			
Minimum doc IPC 7	cumentation searched (classification system followed by classification control to the classifica	ation symbols)	
	tion searched other than minimum documentation to the extent that		
	ata base consulted during the international search (name of data i	base and, where practical, search terms u	sed)
C. DOCUME	ENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with Indication, where appropriate, of the	relevant passages	Relevant to daim No.
A	WO 97 03959 A (WARNER LAMBERT CONTROL OF CHRISTOPHER A (US); JENNINGS RESERVED (1997–02–06) cited in the application		1-6
A	DE 33 27 449 A (GLAXO GROUP LTD 2 February 1984 (1984-02-02) page 9, line 7 - line 16 page 12, line 10 -page 13, line		1-6
A	WO 97 03960 A (WARNER LAMBERT C (US); SCHWEISS DIETER (US)) 6 February 1997 (1997-02-06) cited in the application claim 1	O ;LIN MIN	1-6
		-/	•
X Furt	rther documents are listed in the continuation of box C.	Patent family members are its	sted in annex.
• Special ca	categories of cited documents:	"T" later document published after the	international filling data
	nent defining the general state of the art which is not	or priority date and not in conflict cited to understand the principle of	with the application but
"E" earlier	idered to be of particular relevance r document but published on or after the international	invention "X" document of particular relevance; t	the claimed invention
"L" docum	nent which may throw doubts on priority claim(s) or	cannot be considered novel or ca involve an inventive step when the	nnot be considered to e document is taken alone
O docum	h is cited to establish the publication date of another on or other special reason (as specified) ment referring to an oral disclosure, use, exhibition or	"Y" document of particular relevance; t cannot be considered to involve a document is combined with one of	in inventive step when the or more other such docu-
other	r means nent published prior to the international filing date but than the priority date claimed	ments, such combination being of in the art.	bvious to a person skilled
	man the priority case claimed e actual completion of the international search	"A" document member of the same particle. Date of mailing of the international	
	10 May 2000	17/05/2000	
<u> </u>	d mailing address of the ISA	Authorized officer	
	European Patent Office, P.B. 5818 Patentiaan 2 NL 2280 HV Rijawijk		
1	Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fex: (+31-70) 340-3016	De Jong, B	

1

Intern tal Application No PCT/IB 00/00014

	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 03958 A (WARNER LAMBERT CO ;MCKENZIE ANN T (US)) 6 February 1997 (1997-02-06) cited in the application	1-6
		·

1

information on patent family members

Inten nal Application No
PCT/IB 00/00014

Patent document		Publication		atent family	Publication date
cited in search report		date		nember(s)	date
WO 9703959	A	06-02-1997	AU	6484296 A	18-02-1997
			BG	102187 A	30-10-1998
			BR	9609872 A	23-03-1999
			CA	2220018 A	06-02-1997
			CN	1190955 A	19-08-1998 14-10-1998
			CZ	9800121 A	24-06-1998
			EP	0848705 A	30-04-1998
			HR Hu	960339 A 9900678 A	28-07-1999
			IL	122118 A	14-07-1999
			JP	11509230 T	17-08-1999
		•	NO	980207 A	16-01-1998
			PL	324496 A	25-05-1998
			SK	6298 A	07-10-1998
			US	5969156 A	19-10-1999
					19-10-1999
DE 3327449	Α	02-02-1984	AT	382154 B	26-01-1987
			AT	276783 A	15-06-1986
			AU	566881 B	05-11-1987
			AU	1741783 A	02-02-1984
			BE CA	897422 A 1240313 A	30-01-1984 09-08-1988
			CH	657134 A	15-08-1986
			CZ	280528 B	14-02-1996
			CS	8305687 A	15-03-1988
			CY	1434 A	02-09-1988
			DE	3374010 D	12-11-1987
			DK	68392 A	25-05-1992
			DK	349083 A,B,	31-01-1984
			EP	0107276 A	02-05-1984
			ES.	524590 D	01-06-1985
			ES	8505689 A	01-10-1985
			FI	832757 A,B,	31-01-1984
			FR	2531087 A	03-02-1984
			GB	2127401 A,B	11-04-1984
			GR	79349 A	22-10-1984
			HK	84288 A	28-10-1988
			HÜ	190603 B	29-09-1986
			ΪĒ	55748 B	02-01-1991
			ĪĹ	69375 A	31-12-1986
			IT	1168206 B	20-05-1987
			JP	2025666 C	26-02-1996
			JP	7030084 B	05-04-1995
			JP	59044391 A	12-03-1984
			KE	3805 A	03-06-1988
			KR	9100046 B	19-01-1991
			LU	84935 A	23-11-1983
			MY	5887 A	31-12-1987
			NL	8302705 A	16-02-1984
			NO	832773 A,B,	31-01-1984
			NZ	205083 A	14-03-1986
			PL	243228 A	27-08-1984
			PT	77135 A,B	01-08-1983
			SE	453195 B	18-01-1988
			SE	8304208 A	31-01-1984
			SG	26088 G	15-07-1988
			SI	8311558 A	31-12-1995
			SK	403191 A	11–07–1995

information on patent family members

Inter: nal Application No PCT/IB 00/00014

				1	
Patent document cited in search report		Publication date	Patent famil member(s)		Publication date
DE 3327449	Α		SU 1266	471 A	23-10-1986
			US 4994	567 A	19-02-1991
			US 5013	B33 A	07-05-1991
			US 4562	181 A	31-12-1985
			US 4820	833 A	11-04-1989
				883 A	30-04-1986
				579 A	26-09-1984
			ZW 17	383 A	26-10-1983
WO 9703960	A	06-02-1997	AU 700	794 B	14-01-1999
				896 A	18-02-1997
				188 A	31-08-1998
				714 A	23-02-1999
			CA 2220		06-02-1997
			CN 1190		19-08-1998
				122 A	16-12-1998
				132 A	06-05-1998
				312 A	28-02-1998
				161 A	14-07-1999
			JP 11510		14-09-1999
				209 A	16-01-1998
				463 A	25-05-1998
			SK 5	898 A	05-08-1998
WO 9703958	A	06-02-1997		196 A	18-02-1997
				186 A	30-10-1998
				567 A	06-07-1999 06-02-1997
				458 A	19-08-1998
				1957 A 1123 A	17-06-1998
				1123 A 1704 A	24-06-1998
				313 A	24-00-1998 30-04-1998
				1313 A 1687 A	28-10-1999
				2162 A	14-07-1999
			JP 11509		17-08-1999
				208 A	16-01-1998
				1532 A	08-06-1998
				5998 A	06-05-1998
			31/	MAND W	()()=()¬= : ЧЧА